

Smile Enhancement With Laser Technology—Predictable and Esthetic: A Case Report



by Hugh Flax, D.D.S.

Dr. Flax has been an Accredited Member of the AACD since 1997. He was co-chair of the Conference Advisory Committee for the 2003 Annual Scientific Session and will be for the 2008 meeting in New Orleans. He is a member of the AACD Board of Directors, is on the editorial board of The Journal of Cosmetic Dentistry, and chairs the Disaster Relief Fund. Dr. Flax also is a member of the ADA, the AGD, the ALD, the L.D. Pankey Alumni Association, and the Pierre Fauchard Society. He is a Fellow of the IADFE.

Dr. Flax practices full time in Atlanta, Georgia, focusing on functional- and appearance-related conditions and advanced laser dentistry. He and his wife, Robyn, have two daughters.

INTRODUCTION

With the esthetic zone being absolutely critical to a patient's external appearance and inner emotions, orchestrating a bioesthetic result is mandatory. Too often, this is complicated when esthetic desires infringe on the health of the periodontal complex. This is often true when biologic width violations have occurred iatrogenically. Many factors may contribute to these failures, the two main culprits being intracrevicular margin location and overcontoured restorations. Not only is plaque accumulation problematic, but the supracrestal fibers also become interrupted, causing the tissues to become further inflamed and esthetically unmanageable. Kois' landmark study defined the total dentogingival complex (DGC) as clinically predictable at 3.0 mm on the direct facial aspect, and at 3.0-5.0 mm interproximally when measured from the free gingival margin to the osseous crest. It is critical anteriorly that the gingival margin mimics the osseous scallop while maintaining the DGC.¹ Further complicating these complex situations is the degree of inflammation in the soft tissue, affecting the clinical development of health and esthetic symmetry.

Dental lasers have evolved considerably as an adjunctive and alternative treatment to safely, conservatively, and reliably decrease bacterial levels and improve the hard and soft tissue contours.



Figure 1: Visualizing the entire oral-facial composition helps to diagnose less harmonious features of the smile.

Often the patient is frustrated with his or her previous poor cosmetic results, but to improve the periodontal framework in order to create an ideal result, they must be referred to yet another doctor. Even more challenging is the extended healing time created by reflective mucoperiosteal surgery. This not only affects the chronology of final restorative care, but also delays the patient's ultimate satisfaction and happiness for a minimum of two to three months. Fortunately, dental lasers have evolved considerably as an adjunctive and alternative treatment to safely, conservatively, and reliably decrease bacterial levels and improve the hard and soft tissue contours. Studies of Er: YSGG lasers by Rizoiu and others have shown that thermal coagulative results, as well as bony ablation characteristics are similar to a dental bur.² From a patient-friendly standpoint, less need for suturing and shorter healing times improves case acceptance

for doing ideal dentistry. In selected cases, such as the one presented in this article, minimally invasive laser procedures, with precise restorative planning and technique, can satisfy esthetic and functional parameters. Furthermore, patients can enjoy optimal results more comfortably and efficiently.

A conservative strategy was devised that would allow us to correct the problems and causes in a "multi-tasking" manner.

CASE PRESENTATION

A 38-year-old female patient presented for correction of what she termed her "tilted smile" (Fig 1). Given that she was starting a new sales career, she also wanted to make her teeth brighter and her smile much broader. The patient shared her frustration about previous dental consultations that had

focused solely on orthodontic or surgical solutions without considering a more practical approach that would fit her busy life. Her smile analysis established a collapse of the bicuspid in the buccal corridor. Furthermore, the axial inclinations, irregular gingival margins, and incisal edges created a downward tilt to the patient's right due to tooth positioning. Close-up imaging showed healthy gingival tissues as well as a weakened right central incisor from a large composite (Fig 2).

FINDINGS

A full clinical examination with radiographs and mounted models revealed the following:

- Biomechanically, the majority of her teeth remained strong despite previous dental care.
- Periodontally, soft and hard tissues were healthy.
- Occlusally, load testing was normal (after muscle relaxation)



Figure 2: Close-up photography is essential to planning perio-restorative care.



Figure 3: A mounted diagnostic wax-up is a critical roadmap to planning a realistic result.



Figure 4: Outlining the desired gingival margins, prior to anesthesia, communicates a blueprint to the patient and restorative team.




Figure 5: A stick-bite helps to verify that incisal and gingival planes will be parallel.



Figure 6: The tissues are treated in a very nontraumatic manner with the Waterlase.



Figure 7: To modify the bone, a very tight up-and-down movement is performed, using the black mark as a reference following the gingival scallop.



Figure 8: A curette helps clean and smooth the sulcus of any debris.



Figure 9: A "laser bandage" is placed along the treated area to improve the healing time and decrease the patient's discomfort. Note the immediate improvement of the geometric progression of gingival embrasures.

and there was obvious CR-CO anterior-vertical slide due to a premature contact at tooth #30.

- Esthetically, the width-to-length ratio of the upper centrals was 1:2, far from the ideal range of 0.75:1.0. Tooth shade was a Vita A2.

TREATMENT PLAN

Given the patient's previous history and her desire for minimally invasive dental care, a conservative strategy was devised that would allow us to correct the problems and causes in a "multi-tasking" manner:

- muscle and bite therapy with a Tanner appliance, followed by careful equilibration aided by the T-scan (Tekscan System; South Boston, MA)
- three-dimensional wax-up on a Stratos articulator (Ivoclar Vivadent; Amherst, NY) (Fig 3)
- home bleaching of the lower teeth with Opalescence 15% (Ultradent; South Jordan, UT)
- "closed flap" periodontal modification with the Waterlase ErCr: YSGG (Biolase Technol-

ogy; San Clemente, CA) while the first three items were being accomplished (the combination of these four steps was a tremendous time saver and also allowed us to carefully monitor progress on a weekly basis)

- Lastly, definitive restorative care with porcelain veneers and a crown on tooth #8.

No tissue necrosis or significant bleeding occurred as a result of using the laser's relatively lower settings.

TREATMENT

At the initial closed periodontal lift, the ErCr-YSGG laser was used in three modes (gingival sculpting, osseous recontouring, and bio-stimulation). Prior to anesthesia, the desired framework was planned and outlined using a fine marker (Fig 4). Furthermore, a stick-bite was used, not only to establish an ideal incisal plane, but also to properly align the gingival margins (Fig 5).

With the settings at 2.0 Watts (W), 20 pulses per second, 20% air, and 20% water, a G-6 tip (600 μ in diameter) was used to shape the labial gingival region. No tissue necrosis or significant bleeding occurred as a result of using the laser's relatively lower settings. All areas were "sounded" using a periodontal probe (Fig 6). At the facial margins, osseous sculpting required great precision in order to maintain a 3-mm DGC. A specially tapered T4 tip (400 μ in diameter) was used at a 25% higher wattage of 2.5W. Prior to usage, the tip was measured and marked to 3 mm in order to maintain controlled adjustments within the gingival sulcus during perio probing movement of the tip (Fig 7). The resection was smoothed with a 7/8 curette (Fig 8). Using low-level laser therapy at a setting of 0.25 W, a decrease in the release of inflammatory histamine and increased fibroblasts for junctional epithelial growth was achieved by "frosting" the outer epithelium and injection sites (Fig 9). The patient was placed on a vigorous home-care regimen (Oxygel, Oxyfresh; Coeur d'Alene, ID) and closely monitored



Figure 10: Detailed information helps the laboratory to translate clinical results to the porcelain restorations.



Figure 12: Ideal proportions and emergence profiles will create long-term healthy tissues and bioesthetics.




Figure 11: The great improvement in esthetics boosted the patient's self-confidence and pride in her dental care.

for a month while occlusal therapy and bleaching procedures were performed.

Four weeks after surgery, the tissues had healed and restorative care could be initiated. The patient's teeth were prepared for veneers and a crown with mild soft tissue reshaping, to fine-tune our previous treatment. After taking impressions and bite registrations, prototype provisionals (Luxatemp Plus, Zenith DMG; Englewood, NJ) were fabri-

cated using the "shrink-wrap" technique.

The patient was sent home with the same home-care regimen as mentioned previously, and instructed to "test-drive" her new smile for esthetics and function. She returned in a week to perfect the prototype's occlusion, color, and morphology. Photographs and models were sent to the laboratory, providing a final blueprint for the porcelain restorations (Fig 10).

Four weeks later, the provisionals and cement were carefully removed from the teeth. All restorations were tried in individually and as a group to verify fit and esthetics. After the patient's enthusiastic approval, the porcelain was bonded using the two-by-two technique and isolation. Margins were smoothed and polished and occlusion balanced with the T-scan. A protective nighttime appliance was created to add longevity to the rehabilitation. The very satisfied patient said that we had exceeded her expectations.

CONCLUSIONS

The use of a hard/soft tissue laser is a wonderful adjunctive tool for cosmetic and restorative dentistry. The case discussed here demonstrates that this type of laser technology gives dentists the ability to make significant soft and hard tissue changes while being minimally invasive. These changes not only improve the final esthetic outcome of the case but also provide the physiologic functional parameters required for successful dentistry.

Acknowledgments

The author thanks his office team and laboratory technician, Mr. Wayne Payne (Payne Dental Lab, San Clemente, CA), for continually enhancing the lives of many patients like the one presented here. He also is thankful to his family, who allow him to contribute to the education of other dentists and their teams.

References

1. Kois JC. Altering gingival levels: The restorative connection, Part I: Biologic variables. *J Esthet Dent* 6(1):3-9, 1994.
2. Rizoiu I, et al. Osseous repair subsequent to surgery with an erbium hydrokinetic laser system (pp.213-221). International Laser Congress, International Proceedings Division. Athens, Greece, September 25-28, 1996. *Ap*

